

*Erratum*

**Erratum to ‘Spasticity and Range of Motion Over Time in Stroke Patients Who Received Multiple-Dose Botulinum Toxin Therapy’ [Journal of Stroke and Cerebrovascular Diseases, Vol. 29, No. 1 (January), 2020: 104481]**

Takanori Ro, RPT,\* Tetsuo Ota, MD, PhD,† Tsukasa Saito, MD, PhD,‡ and  
Ou Oikawa, MD, PhD†

---

The publisher regrets that an error in [Table 2](#) was included in the original published article. The correct version of [Table 2](#) is included below. The publisher would like to apologise for any inconvenience caused.

---

From the \*Rehabilitation Unit, Asahikawa Medical University Hospital, Asahikawa, Japan; †Department of Physical Medicine and Rehabilitation, Asahikawa Medical University Hospital, Asahikawa, Japan; and ‡Cardiovascular, Respiratory and Neurology Division, Department of Internal Medicine, Asahikawa Medical University, Asahikawa, Japan.

Address correspondence to Takanori Ro, RPT, Rehabilitation Unit, Asahikawa Medical University Hospital, 2-1-1-1, Midorigaoka-Higashi, Asahikawa, Hokkaido 078-8510, Japan. E-mail: [keio\\_tukigase\\_ro@yahoo.co.jp](mailto:keio_tukigase_ro@yahoo.co.jp).

DOI of original article: <http://dx.doi.org/10.1016/j.jstrokecerebrovasdis.2019.104481>.

1052-3057/\$ - see front matter

© 2020 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.104769>

**Table 2.** *The number of units of BoNT-A injection and the percentage of patients injected*

First dose (24 patients)	Mean (U)	SD	Percentage (%)
Total	272.6	82.1	
Greater pectoral	42.5	10.6	8.7
Biceps	61.4	21.9	47.8
Brachioradialis	60.0	0.0	4.3
Flexor carpi radialis	42.3	10.8	47.8
Flexor carpi ulnaris	42.5	9.2	43.5
Flexor digitorum superficialis	41.8	12.7	60.9
Flexor digitorum profundus	39.2	12.6	56.5
Flexor pollicis longus	18.0	4.5	21.7
Adductor muscle of the thumb	20.0	0.0	13.0
Lateral gastrocnemius	52.5	19.4	52.2
Medial gastrocnemius	52.5	19.4	52.2
Soleus	58.1	17.1	34.8
Tibialis posterior	60.0	14.7	60.9
Flexor digitorum longus	52.5	20.8	52.2
Flexor hallucis longus	60.0	0.0	8.7
Second dose (24 patients)	Mean (U)	SD	Percentage (%)
Total	302.6	63.8	
Biceps	54.2	18.7	52.2
Brachioradialis	44.5	20.5	8.7
Flexor carpi radialis	33.2	12.2	60.9
Flexor carpi ulnaris	33.9	12.9	60.9
Flexor digitorum superficialis	33.2	11.7	73.9
Flexor digitorum profundus	35.3	13.0	78.3
Flexor pollicis longus	21.1	7.8	39.1
Extensor hallucis longus	30.0	0.0	4.3
Adductor muscle of the thumb	12.5	10.6	8.7
Lateral gastrocnemius	59.3	12.2	65.2
Medial gastrocnemius	59.3	12.2	65.2
Soleus	58.8	7.9	34.8
Tibialis posterior	57.5	14.8	69.6
Flexor digitorum longus	48.5	15.6	43.5
Flexor hallucis longus	45.0	12.9	17.4
Third dose (24 patients)	Mean (U)	SD	Percentage (%)
Total	303.9	67.7	
Biceps	52.5	25.6	52.2
Triceps	60.0	28.3	8.7
Brachioradialis	45.0	12.9	17.4
Flexor carpi radialis	32.5	10.8	69.6
Flexor carpi ulnaris	32.5	10.8	69.6
Flexor digitorum superficialis	34.2	10.2	82.6
Flexor digitorum profundus	33.7	10.7	82.6
Flexor pollicis longus	14.3	5.3	30.4
Extensor hallucis longus	27.5	3.5	8.7
Adductor muscle of the thumb	16.7	5.8	13.0
Lateral gastrocnemius	50.7	15.3	60.9
Medial gastrocnemius	52.9	11.6	60.9
Soleus	54.2	13.6	26.1
Tibialis posterior	51.2	11.8	73.9
Flexor digitorum longus	47.0	15.8	65.2
Flexor hallucis longus	43.8	14.9	17.4

Fourth dose (24 patients)	Mean (U)	SD	Percentage (%)
Total	326.3	55.3	
Biceps	44.3	11.8	65.2
Triceps	35.0	7.1	8.7
Brachioradialis	47.0	8.4	21.7
Flexor carpi radialis	29.7	12.7	69.6
Flexor carpi ulnaris	29.7	12.7	69.6
Flexor digitorum superficialis	29.5	12.0	91.3
Flexor digitorum profundus	27.9	11.3	82.6
Flexor pollicis longus	16.7	5.2	26.1
Extensor hallucis longus	25.0	7.1	8.7
Adductor muscle of the thumb	10.0	0.0	8.7
Lateral gastrocnemius	54.4	16.4	78.3
Medial gastrocnemius	54.4	16.4	78.3
Soleus	64.2	9.2	26.1
Tibialis posterior	50.3	7.8	78.3
Flexor digitorum longus	51.6	8.9	69.6
Flexor hallucis longus	38.0	13.0	21.7
Fifth dose (24 patients)	Mean (U)	SD	Percentage (%)
Total	339.1	33.9	
Greater pectoral	50.0	0.0	9.1
Biceps	47.8	10.9	40.9
Triceps	40.0	10.0	13.6
Brachioradialis	50.0	0.0	13.6
Flexor carpi radialis	36.5	9.9	59.1
Flexor carpi ulnaris	36.5	9.9	59.1
Flexor digitorum superficialis	39.0	8.4	90.9
Flexor digitorum profundus	40.0	7.3	86.4
Flexor pollicis longus	21.1	9.3	40.9
Extensor hallucis longus	20.0	0.0	4.5
Adductor muscle of the thumb	20.0	0.0	4.5
Lumbricales	20.0	0.0	4.5
Lateral gastrocnemius	49.1	14.9	72.7
Medial gastrocnemius	49.1	14.9	72.7
Soleus	52.9	15.0	31.8
Tibialis posterior	54.2	8.4	86.4
Flexor digitorum longus	52.9	6.1	63.6
Flexor hallucis longus	42.0	13.0	22.7